

Constraints on RRAT Emission Mechanisms from *RXTE*/PCA Observations of RRAT J1819–1458

Robert E. Rutledge¹

ABSTRACT

We derive the second and most stringent limit to date of the X-ray/radio flux ratio ($F_X/F_{1.4\text{GHz}}$) for the radio bursts associated with the recently identified source class, the Rotating Radio Transients (RRATs). We analyze 20.1 hr of *RXTE*/PCA observations of RRAT J1819–1458 – a period during which 350 ± 23 RRAT radio bursts occurred, based on the previously observed average radio burst rate. No X-ray bursts were detected, implying an upper-limit on the X-ray flux for RRAT-bursts of $F_{X,\text{lim}} < 1.5 \times 10^{-8} \text{ erg cm}^{-2} \text{ s}^{-1}$ (2–10 keV), or a luminosity $< 2.3 \times 10^{37} (d/3.6 \text{ kpc})^2 \text{ erg s}^{-1}$. The time-average burst flux is $< 2 \times 10^{-13} \text{ erg cm}^{-2} \text{ s}^{-1}$ (0.5–8 keV) – a factor of 10 below that of the previously identified persistent X-ray counterpart. Thus, X-ray bursts from the RRAT are energetically unimportant compared with the persistent X-ray emission. From the previously observed burst radio flux, we derive an upper-limit $F_X/F_{1.4\text{GHz}} \leq 4.2 \times 10^{-12} \text{ erg cm}^{-2} \text{ s}^{-1} \text{ mJy}^{-1}$ for the radio bursts from this RRAT, the most stringent to date, due to the high radio flux of bursts from this source. The $F_X/F_{1.4\text{GHz}}$ ratio is a factor ~ 80 larger than that of the millisecond pulsar PSR B1821–24; thus emission processes of X-ray/radio efficiency comparable to MSP pulses cannot be ruled out. However, if the RRAT burst emission mechanism is identical to the ms bursts of magnetars, then the ms bursts of magnetars should be easily detected with radio instrumentation; yet none have been reported to date.

Subject headings: stars: flare; stars: neutron; X-rays: stars; pulsars: individual (J1819–1458)

1. Introduction

Recently, a new observational class of neutron stars has been proposed (McLaughlin et al. 2006, M06 hereafter), Comprised of eleven sources, the class is defined by the phe-

¹Department of Physics, McGill University, 3600 rue University, Montreal, QC, H3A 2T8, Canada; rutledge@physics.mcgill.ca

nomenon of fast (2-30 ms) radio bursts, which repeat on timescales of 4 min - 3 hr. Periodicities were claimed to be found in the range of $P = 0.4-7$ s for ten of the eleven sources, through period folding; this suggested association with rotating neutron stars. In three of the eleven sources, period derivatives were claimed, with one suggesting a magnetic field strength of 5×10^{13} G – a possible magnetar. Because the $P - \dot{P}$ analysis remains unpublished and undescribed, we regard association of the radio-bursting population with neutron stars to be an unsupported hypothesis; nonetheless, we adopt the hypothesis and nomenclature of referring to the sources as Rotating RAdio Transients (RRATs), and consider their association with neutron stars to be the leading hypothesis.

One of these – RRAT J1819–1458 – has a $7'' \times 32''$ error ellipse (M06)¹, which permitted discovery of an X-ray counterpart, CXOU J181934.1–145804 (Reynolds et al. 2006), with absorption-corrected flux of 2×10^{-12} erg cm⁻² s⁻¹ (0.5-8 keV), or 3.1×10^{33} erg s⁻¹ at the $d = 3.6$ kpc dispersion-measure-indicated distance. Two-hundred twenty nine radio bursts at $\nu = 1.4$ GHz, with average half-duration $w_{50} = 3$ ms were observed in 13 hours of radio integration, the brightest with a peak flux of $S_\nu = 3.6$ Jy. If we take w_{50} to be the light crossing time of the emission region, this implies a source brightness temperature (e.g., Güdel 2002):

$$T_b = 3 \times 10^{23} \left(\frac{S_\nu}{3.6 \text{ Jy}} \right) \left(\frac{1.4 \text{ GHz}}{\nu} \right)^2 \left(\frac{3 \text{ ms}}{w_{50}} \right)^2 \left(\frac{d}{3.6 \text{ kpc}} \right)^2 \text{ K} \quad (1)$$

This is significantly above the brightness temperature limit for inverse Compton scattering ($T_B \sim 10^{12}$ K) which likely requires a coherent emission mechanism for this and other RRATs, possibly accompanied by X-ray emission.

In this paper, we analyze *RXTE*/PCA data, and search for X-ray bursts which may accompany the observed radio bursts, finding none. In § 2, we describe the observations and analysis; in § 3, we present the results of the X-ray burst search; and in § 4, we discuss the implications of these results and conclude.

¹It is unclear why the positional uncertainty from this source is so much smaller than, for example, the $15' \times 7'$ error region for RRAT J1911+00, obtained in the same study with the same instrumentation.

2. Observations and Analysis

We analyze observations taken in *RXTE*/PCA (Swank et al. 1996; Jahoda et al. 2006), obtained from the public archive (Table 1). For basic data extraction, we used FTOOLS² (Blackburn 1995) following standard *RXTE* recipes. We used `GoodXenon_16s`-type data, analyzing 20.1 hr of data. We performed some data selections: we excluded data from PCU1 (Proportional Counter Unit 1, numbered 0-4), and layers 2-3 from all PCUs, and required at least 2 active PCUs. Most observations subsequently had 3 or 4 active PCUs (0,2-4; or 0,2-3). These selections eliminate the PCU high-voltage breakdowns³ which occur in PCU1, PCU3 and PCU4 (Jahoda et al. 2006) in these and other data on which the programs were tested.

We examined pulse-height-analyzer (PHA) channels only ≤ 27 corresponding to an approximate energy range of 1.5-10 keV, and excluded propane-veto layer events.

We searched for bursts on two timescales: $\tau = 3.0$ ms (the observed radio-burst timescale for RRAT J1819–1458) and $1/10$ that value (0.3 ms), to cover possible shorter X-ray duty cycles.

We measure the number of counts M over a period of ± 1 sec about each count in the data stream. We measure the number N of counts which are observed within $\pm \tau/2$ of each count, and we calculate the Poisson probability of measuring $\geq N$ counts for an expected number of counts $\mu = M \times \tau/(2 \text{ s})$ as:

$$P_{\text{Poisson}}(\mu, \geq N) = \sum_{i=N}^{\infty} \frac{\mu^i}{i!} \exp(-\mu)$$

Summing over all 22 observations, we observed $N_{\text{phot}} = 3069964$ total counts. An event for which $P_{\text{Poisson}} \times N_{\text{phot}} < 0.01$ would have 1% chance of being produced due to random fluctuation alone during the entirety of our analysis; we call such an event a “trigger”, and store all information about the $N + 1$ counts in that trigger for future examination.

²<http://heasarc.gsfc.nasa.gov/ftools>

³HV breakdown in PCU3 and PCU4 is typically marked by a burst of counts, usually in PHA channels < 20 , and limited to a single wire within the PCU, on a ≤ 2 ms timescale. However, breakdown in PCU1 is marked by low-PHA counts across *all* wires, and is often terminated by a single high-energy (PHA 15-30) count; it may be due to a distinct phenomenon.

3. Results

We found no triggers in the above data stream, on either timescale. We calculated the limits on burst rate as a function of average burst flux over the 3 ms and 0.3 ms timescale (Fig. 1). At the time of each count, we calculated the average (over τ) number of counts per PCU per sec, corresponding to a 99% confidence countrate fluctuation (that is, a Poisson fluctuation which would only occur, on average, in a dataset with $100\times$ as many counts as the present dataset); this resulted in 99% confidence upper-limits between 422-1010 c/s/PCU on the $\tau = 3\text{ms}$ timescale). During these observations, RRAT J1819–1458 lay $27.5'$ off-axis, at a collimator efficiency of 0.45-0.5, for which we correct the countrates. Layer-1 wires in the PCUs make up $\approx 90\%$ of the total active area below 10 keV, for which we correct the countrate (the remaining 10% is in layers 2-3, which we neglected). Finally, we use an energy correction factor of 1.0×10^{-8} erg $\text{cm}^{-2} \text{s}^{-1}$ (2-10 keV) per 1000 c/s/PCU (assuming a photon power-law slope $\alpha=1$, and neglecting N_{H}^4 which we assume as the default spectrum throughout), where here, the countrates are for layers 1-3. The maximum burst rate is the 95% confidence upper-limit for the burst rate, $2./T(> F_{\text{lim}})$, where $T(> F_{\text{lim}})$ is the total integrated observing time sensitive to bursts from RRAT J1819–1458 with flux $> F_{\text{lim}}$; the numerator (2) comes from the number of bursts for which one detects 0 bursts $\leq 5\%$ of the time.

In general, flux sensitivity did not change much during observations; thus, we detected no bursts on a $\tau = 3\text{ms}$ ($\tau = 0.3\text{ms}$) timescale at fluxes above $\gtrsim 1.5\times 10^{-8}$ erg $\text{cm}^{-2} \text{s}^{-1}$ (8×10^{-8} erg $\text{cm}^{-2} \text{s}^{-1}$), with 99% confidence, with burst rates as low as 0.1hr^{-1} . At lower fluxes, the observations rapidly become insensitive.

4. Discussion and Conclusions

RRAT J1819–1458 is the most prolific radio-burster of all the reported RRATs – 229 radio bursts detected in 13 hr of radio observations. On average, we would expect 350 ± 23 such radio bursts to have occurred during these observations, assuming the radio burst rates do not change with time. None of these had a corresponding X-ray flux $\geq 1.5\times 10^{-8}$ erg $\text{cm}^{-2} \text{s}^{-1}$ (2-10 keV). We expect 2, on average, to have been as radio bright as the brightest previously observed (3.6 Jy), which implies that ≥ 1 such bursts were emitted (95% confidence). This, in combination with our derived burst flux limit, results in an X-ray to radio flux ratio limit for bursts from RRAT J1819–1458 of approximately:

⁴using WebPIMMS; <http://heasarc.gsfc.nasa.gov/Tools/w3pimms.html>

$$F_X/F_{1.4\text{ GHz}} \leq 4.2 \times 10^{-12} \text{ erg cm}^{-2} \text{ s}^{-1} \text{ mJy}^{-1}$$

This limit is a factor of 14 lower than that derived for RRAT J1911+00 Hoffman et al. (2006), due to the factor of 14 greater radio flux observed from the brightest radio burst from RRAT J1819–1458 (M06). It may be instructive to compare this ratio with that from other transient coherent emission processes.

The limit corresponds to $F_X/F_{1.4\text{ GHz}} \leq 10^{14.7}$ Hz, comparable to the average ratio observed from stellar coronal activity (Guedel & Benz 1993); however the timescales for coronal flares (for example) are typically $\sim 10^3$ sec, much longer than the ms flares of RRATs.

Compared to MSPs, this upper-limit to $F_X/F_{1.4\text{ GHz}}$ is a factor 84 above the observed ratio for (for example) the millisecond pulsar PSR B1821-24 ($F_X/F_{1.4\text{ GHz}} = 5 \times 10^{-14} \text{ erg cm}^{-2} \text{ s}^{-1} \text{ mJy}^{-1}$; Hoffman et al. 2006). Thus, this limit does not exclude an emission mechanism for RRAT bursts with comparable X-ray to radio efficiency to MSPs emission mechanism.

The claim of measured $P - \dot{P}$ values (M06) which imply magnetic field strengths approaching those of magnetars, makes comparison between the RRAT-bursts and those from magnetars a natural one, with the expectation that the emission mechanisms may be similar. The absence of X-ray/radio burst flux limits for the ms X-ray bursts of magnetars (Göğüş et al. 1999; Göğüş et al. 2001; Gavril et al. 2002; Kaspi et al. 2003; Gavril et al. 2004; Woods et al. 2005) precludes direct comparison with the emission mechanism powering those sources. However, a radio observation of such would be valuable to compare with the derived $F_X/F_{1.4\text{ GHz}}$ limits for RRATs. Taking 1E 2259+586 as an example (Gavril et al. 2004), with median duration of ~ 100 ms, and peak fluences of $10^{-9} \text{ erg cm}^{-2}$, magnetar bursts would produce coincident radio bursts of $\geq 2500 \text{ mJy}$, if the magnetar bursts were due to a similar emission mechanism as RRATs. This radio flux is comparable to the brightest RRAT radio bursts yet observed (M06), and are therefore readily detectable, yet no such bursts have been reported. However, the burst rate of the ms X-ray bursts of magnetars is not constant in time; whether this is true for the radio bursts of RRATs has not been described in the reported observations (M06). On the other hand, a non-detection of radio bursts from magnetars to flux limits below this value would require a different $F_X/F_{1.4\text{ GHz}}$ ratio for the ms bursts from magnetars, and may therefore imply a distinct emission mechanism.

A burst flux limit $F_{\text{lim}} < 1.5 \times 10^{-8} \text{ erg cm}^{-2} \text{ s}^{-1}$ (2-10 keV), averaged over 3 ms, combined with an average burst rate of 228/13 hr (M06), results in time-average burst flux of $< 2.2 \times 10^{-13} \text{ erg cm}^{-2} \text{ s}^{-1}$ (2-10 keV), or, extrapolating to the energy band for the X-ray counterpart (Reynolds et al. 2006) $< 2.1 \times 10^{-13} \text{ erg cm}^{-2} \text{ s}^{-1}$ (0.5-8 keV). The time-average burst X-ray flux is a factor of 10 less than the persistent X-ray counterpart in this pass-band

Reynolds et al. (2006). Thus, X-ray bursts related to the radio bursts of this RRAT are energetically unimportant compared to the persistent X-ray flux. In addition, the time-average burst X-ray efficiency $\eta = L_X(0.5 - 8 \text{ keV})/\dot{E}$, compared with the reported dipolar emission $\dot{E} = 24.94(5) \times 10^{31} \text{ erg s}^{-1}$ (M06) is $\eta \leq 7 \times 10^{-3} (d/3.6 \text{ kpc})^2$.

It is unlikely that great improvement will be made in the X-ray flux limit presented here using existing instrumentation. An improved flux limit cannot be obtained with *Chandra*: 2 counts in the HRC-S detected within 2 ms corresponds to $3 \times 10^{-8} \text{ erg cm}^{-2} \text{ s}^{-1}$ (extrapolated to 2-10 keV). On-source *RXTE*/PCA observations could improve on the present flux sensitivity by a factor of $\times 2$.

RER acknowledges support from NSERC through the Discovery Grant program. RER acknowledges useful conversations with Marjorie Gonzalez, Margaret Livingstone, Vicky Kaspi and Andrew Cumming.

REFERENCES

- Blackburn, J. K., 1995, in R. A. Shaw, H. E. Payne, & J. J. E. Hayes (eds.), *ASP Conf. Ser. 77: Astronomical Data Analysis Software and Systems IV*, p. 367
- Gavriil, F. P., Kaspi, V. M., & Woods, P. M., 2002, *Nature* 419, 142
- Gavriil, F. P., Kaspi, V. M., & Woods, P. M., 2004, *ApJ* 607, 959
- Gögüş, E., Woods, P. M., Kouveliotou, C., van Paradijs, J., Briggs, M. S., Duncan, R. C., & Thompson, C., 1999, *ApJ* 526, L93
- Gögüş, E., Kouveliotou, C., Woods, P. M., Thompson, C., Duncan, R. C., & Briggs, M. S., 2001, *ApJ* 558, 228
- Güdel, M., 2002, *ARA&A* 40, 217
- Guedel, M. & Benz, A. O., 1993, *ApJ* 405, L63
- Hoffman, K., Rutledge, R. E., Fox, D. B., Gal-Yam, A., & Cenko, S. B., 2006, *ApJ*, submitted, astro-ph/0609092
- Jahoda, K., Markwardt, C. B., Radeva, Y., Rots, A. H., Stark, M. J., Swank, J. H., Strohmayer, T. E., & Zhang, W., 2006, *ApJS* 163, 401

- Kaspi, V. M., Gavriil, F. P., Woods, P. M., Jensen, J. B., Roberts, M. S. E., & Chakrabarty, D., 2003, *ApJ* 588, L93
- McLaughlin, M. A., Lyne, A. G., Lorimer, D. R., Kramer, M., Faulkner, A. J., Manchester, R. N., Cordes, J. M., Camilo, F., Possenti, A., Stairs, I. H., Hobbs, G., D’Amico, N., Burgay, M., & O’Brien, J. T., 2006, *Nature* 439, 817
- Reynolds, S. P., Borkowski, K. J., Gaensler, B. M., Rea, N., McLaughlin, M., Possenti, A., Israel, G., Burgay, M., Camilo, F., Chatterjee, S., Kramer, M., Lyne, A., & Stairs, I., 2006, *ApJ* 639, L71
- Swank, J. H., Jahoda, K., Zhang, W., & Giles, A. B., 1996, in M. A. Alpar, U. Kiziloglu, & J. Van Paradijs (eds.), *The Lives of the Neutron Stars*, NATO ASI Ser C., 450), Boston: Kluwer
- Woods, P. M., Kouveliotou, C., Gavriil, F. P., Kaspi, V. M., Roberts, M. S. E., Ibrahim, A., Markwardt, C. B., Swank, J. H., & Finger, M. H., 2005, *ApJ* 629, 985

Fig. 1.— Upper-limits on burst rate as a function of flux, averaged on a 3 ms (solid line) and 0.3 ms timescales, from RRAT J1819–1458 (see § 3 for discussion).




Figure 1

Table 1. *RXTE* Observations of RRAT J1819–1458

Start Time (TT) (dd/mm/yy hh:mm)	End Time (TT) (dd/mm/yy hh:mm)	(ksec)	# PCUs ^b	N_{phot}	$\langle I \rangle$ (c/s) ^a (1.5-10 keV)
06/06/98 12:50	06/06/98 13:42	3.16	4	155048	49.07
06/06/98 14:25	06/06/98 15:25	3.63	4	183683	50.57
06/06/98 16:02	06/06/98 17:01	3.53	4	182342	51.57
06/06/98 17:42	06/06/98 18:37	3.31	4	159157	48.05
08/06/98 09:39	08/06/98 10:25	2.73	4	124533	45.52
08/06/98 11:15	08/06/98 12:01	2.73	4	119602	43.71
08/06/98 12:51	08/06/98 13:43	3.13	4	143406	45.73
08/06/98 14:27	08/06/98 15:26	3.55	4	167412	47.13
08/06/98 16:03	08/06/98 17:03	3.61	4	172435	47.69
08/06/98 17:43	08/06/98 18:40	3.39	4	161687	47.67
10/06/98 12:54	10/06/98 13:45	3.02	4	129359	42.78
10/06/98 14:28	10/06/98 15:27	3.56	4	158155	44.33
10/06/98 16:04	10/06/98 17:05	3.64	4	158619	43.48
10/06/98 19:21	10/06/98 20:17	3.34	2	140593	42.06
10/06/98 20:58	10/06/98 21:53	3.26	4	131917	40.42
10/06/98 22:34	10/06/98 23:14	2.43	4	97358	40.03
12/06/98 11:18	12/06/98 12:02	2.67	3	98495	36.88
12/06/98 12:53	12/06/98 13:46	3.21	4	134874	41.94
12/06/98 14:29	12/06/98 15:28	3.55	4	120253	33.86
12/06/98 16:05	12/06/98 17:05	3.61	3	113601	31.42
12/06/98 17:41	12/06/98 18:41	3.63	3	111473	30.69
12/06/98 19:19	12/06/98 20:17	3.50	3	105962	30.24

^aAverage countrate during observation for PHA channels ≤ 27 .

^bNumber of active PCUs at observation start time. PCUs turn on and off during the course of the observations.